



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Ecological and Biological Aspects of Pasture Management in Mongolia

Tserendash Sainkhoo

Research Institute of Animal Husbandry, Mongolia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/18-1/27>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Ecological and biological aspects of pasture management in Mongolia

Tserendash Sainkhuu , Sc.D ,Professor Director ,Research Institute of Animal Husbandry
Zaisan-210153 , Ulaanbaatar ,Mongolia . E-mail :*riakh@maginet.mn ; tsdash@yahoo.com*

Key words : pasture management , productivity , carrying capacity , rotation

Introductions Mongolia has 147 .1 million ha area suitable for pasture , which is a basic fodder for animal husbandry of Mongolia . Pasture management is only way of prevention degradation and year round availability of feed for livestock sector will be provided due to the pasture management . Temporally , the pasture management is divided into 2 periods such as warm and cold season . Warm season pasture management requires the most responsible and accurate actions . During warm period or 120 days , plant growth and development undergoes numbers of stages , and achievement to definite level of productivity , followed by withering takes place . Main goal of pasture management is provide the condition to complete all growth stages of plants and define grazing duration . Most sensitive growth period which is to define further growth of the plants is tillering , budding and heading . If livestock grazing coupled with plant tillering , shooting then decrease productivity totally . Therefore , it would be important to manage pasture utilization to avoid overgrazing in first stages of pasture plants .

Materiers and methods Experimental data gathered during field study of rangelands encompassing 4 ecological zones of Mongolia in 1972-1990 and 1991-2005 are the basis of the paper . These experimental data include more than 1500 descriptions of soil and plant communities , more than 22000 samples of vegetation , plant species collected for yield . Basical coverage of plant species , rocks , litter and bareground were determined in randomly selected areas . The productivity of pastures was measured in May-August , in October , in March and in April (S.Tserendash ,1972) . Plant samples were classified as green grass and litter . Carrying capacity and stocking rate of pasture were determined by S.Tserendash (1996) .

Results and conclusions The accumulation of pasture productivity is variable in different regions depending on the species compositions , bio morphological characteristics , ecological and climatic conditions . The maximal productivity of any pastures is found during end of flowering and seeding stages of the plants , dominant for late July or first decade of August , and biological productivity is 10 .5-15 .0 c/ha in high mountainous region , 11 .5-19 .4 c/ha in forest-steppe , 6 .5-13 .0 c/ha in steppe and 2 .9-3 .8 c/ha in semi-arid regions . There is a carrying capacity to bear in an average 86 million sheep in normal climate condition per year . It is necessary to estimate the pasture grazing capacity in our country in 3 versions , such as drought years , normal year's and favorable year's by relying on the multi-year changes of vegetation . Research result demonstrates that the use of forest steppe , steppe and semi-arid pastures in tillering stage or late May is activated , resulted in 2-4 fold regrowth as multi-year average . It is therefore possible to reuse the forest-steppe pastures twice or thrice , steppe pasture once or twice and high mountain pasture once relied on the characteristics of vegetation regrowth . The above mentioned principle on vegetation regrowth of various pastures is important for estimation of the adequacy , grazing and stocking capacities of summer pastures , and it is a source for reserve . Use of pasture aftermath provides with the possibility of using 10-40% greater protein and 10-30% greater yield per unit of area .

Any pasture is composed of numbers of plant species growing in harmony as a result of evolution . Of these plant species only 3-5 species of plants drive the pasture growth and development . As well as the productivity and stability of any pasture types depends on these plants . Grass is advanced then others by their function in vegetation cover , climate , tolerance human activity impact , competitive capacity . Pasture sustainable management depend on biology , morphology , and abundance of functional group of plants . Total pasture productivity can be determined by grass yield based grass steady growth as well as pasture improvement technology may proceed . If grass makes up 25 % pasture productivity , it would be give positive result to shift pasture plant growth periods .

Proper combination of the duration and frequency of pasture use with patterns of plant development is a basis of organizing pasture uses as well as protecting it from degradation . The best way to adapt pasture vegetation for climate change is to follow a concept to use 50 % of aboveground grass biomass and leave 50% on the ground .